

For use around buildings and structures for the control of listed pests, including on lawns, around the base of ornamental trees and shrubs, around residential, institutional, public, commercial, agricultural and industrial buildings; and on parks, on recreational areas, and on athletic fields.

GUARANTEED ANALYSIS	
Total Nitrogen (N) .....	17.00%
17.00% Urea Nitrogen*	
Soluble Potash .....	3.00%

Information regarding the contents and level of metals in this product is available on the Internet at <http://www.aapfco.org/metals.html>

Imidacloprid .....	0.20%
Lambda-cyhalothrin .....	0.04%
OTHER INGREDIENTS .....	99.76%
TOTAL: .....	100.00%

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION/PRECAUCION**  
Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand this label, find someone to explain it to you in detail.)

FIRST AID	
<b>IF IN EYES</b>	<ul style="list-style-type: none"> <li>Hold eye open and rinse slowly and gently with water for 15 to 20 minutes</li> <li>Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
<p align="center"><b>HOT LINE NUMBER</b></p> <p>In case of medical emergency only call (877) 325-1840. Have a product container or label with you when calling a poison control center or doctor for treatment advice.</p>	
<p align="center"><b>NOT TO PHYSICIAN</b></p> <p>No specific antidote is available. Treat the patient symptomatically.</p>	

**PRECAUTIONARY STATEMENTS**  
**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**  
**CAUTION**

**ENVIRONMENTAL HAZARDS**

This product is highly toxic to fish and aquatic invertebrates. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate. Runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

This product contains a chemical with properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

Users Should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**Use Restrictions:**

- Keep children and pets off treated area until product has been watered in and lawn is dry.
- Do not apply this product to established growing crops or stored raw agricultural commodities used for food or feed.
- Do not allow applications to contact water inhabited by fish, such as in ornamental fish ponds that are located around structures being treated.
- Do not apply this product by aerial application.
- Apply only to actively growing turf. Do not apply near water, storm drains, or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn, and sweep any product that lands in the driveway, sidewalk, or street, back onto your lawn. Check with your local Cooperative Extension Agency to obtain specific information on local turf best management practices. Check with your county or city government to determine if there are local regulations for fertilizer use.

- Do not apply more than 4.6 lb of this product per 1,000 square feet or 200 lb of product (0.4 lb of imidacloprid active ingredient) per acre per year.
- Not for use in commercial greenhouses, nurseries, on sod farms or on grass grown for seed

Consult your local State Agricultural Experiment Station or State Extension Turf Specialists for more specific information regarding timing of application.

<b>RATES OF THIS PRODUCT FOR GRUB AND SOIL INSECT CONTROL ON TURFGRASS</b>		
<b>PEST</b>	<b>USE DIRECTIONS</b>	<b>RATE</b>
<b>Larvae of:</b> European Crane Fly, ( <i>Tipula paludosa</i> ) Annual bluegrass weevil, ( <i>Hyperodes</i> spp.) Green June Beetle, ( <i>Cotinis nitida</i> ); May or June beetle, ( <i>Phyllophaga</i> spp.); Asiatic garden beetle, ( <i>Maladera castanea</i> ) Japanese Beetle, ( <i>Popillia japonica</i> ) Billbug, ( <i>Sphenophorus</i> spp.) Northern masked chafer and Southern masked chafer, ( <i>Cyclocephala borealis</i> , <i>C. immaculata</i> , and/or <i>C. lurida</i> ) Black turfgrass ataenius, ( <i>Ataenius spretulus</i> and <i>Aphodius</i> spp.) Oriental Beetle, ( <i>Anomala orientalis</i> ) Cutworms (suppression), European chafer, ( <i>Rhizotrogus majalis</i> )	For control of grubs, billbugs, european crane fly and annual bluegrass weevil, make application prior to egg hatch of the target pest.	3.4 - 4.6 lb per 1,000 sq ft  (150 - 200 lbs per acre)
Chinchbug Mole Crickets, ( <i>Scapteriscus</i> spp.)	For control of chinchbugs, make application prior to hatching of the instar nymphs.  For control of mole crickets make application prior to or during the peak egg hatch period. When adults or large nymphs are present and actively tunneling, apply with a remedial insecticide.	4.6 lb per 1,000 sq ft  (200 lbs per acre)

Use the higher listed rate for longer length of control. Watering (irrigation or rainfall) following application will dissolve the granules and release the insecticide. Uniform coverage is necessary to provide the most effective level of control. Consult your state university or local Cooperative Extension Service office for specific pest control application timing in your area. Start application to turf prior to the establishment of high insect pest populations and significant turf damage. Make reapplications as necessary to keep pest populations under control, using higher rates as pest pressure increases.

TREATMENT	PEST	Rate of product per 1000 ft <sup>2</sup>	Rate of Product per Acre
<b>Band Treatment Along Foundation</b> (not to exceed 3 ft from foundation; only over soil or plants)	Ants (Southern, Field, Allegheny Mound, Cornfield, Honey, Pavement, Nuisance and Fire Ants) (excluding carpenter and pharaoh) Beetles Crickets Earwigs Firebrats Fleas Millipedes Mites Ticks	2.2 – 3.3 lbs.	97 – 144 lbs.
<b>Lawn Application</b>	Ants (Southern, Field, Allegheny Mound, Cornfield, Honey, Pavement, Nuisance and Fire ants) (excluding carpenter and pharaoh) Armyworms (Fall & Yellowstriped) Crickets Cutworms Earwigs European Crane Flies Fleas Grasshoppers Green June Beetles Japanese Beetle (Adult) Leafhoppers Millipedes Mites Lawn moths (Sod webworms) Spider Mites (Twospotted, Spruce) Spittlebugs Springtails Ticks	2.2 – 3.3 lbs.	97 – 144 lbs.

<b>Lawn Application</b>	Billbugs Chinch Bugs Dung beetles* Green June beetle grubs* Hyperodes weevils Japanese beetle grubs* Mole Crickets	3.3 – 4.4 lbs	144 – 194 lbs.
<b>Spot Treatment</b> (Not for Use in California)	Ants (excluding carpenter and pharaoh)	Apply 1 teaspoon to ½ cup over, and 2 ft. around each ant mound	
* Suppression only			

PEST	RATE	USE INSTRUCTIONS
Ants (Southern, Field, Allegheny Mound, Cornfield, Honey, Pavement, Nuisance and Fire Ants) (excluding carpenter and pharaoh)	Apply at the rate of 2.2 lb to 3.3 lb per 1,000 sq. ft. (an area 5' x 200') oz.	Apply up to a 3 foot band of granules next to the house foundation and completely distribute the granules. Water lightly immediately following application.
Beetles	or	Treat when insects first appear or when you notice lawn damage.
Crickets		
Earwigs	3.5 oz. to 5.3 oz. per 100 sq. ft. (10' x 10').	Remove leaf litter and debris from the foundation area. Cut back branches and vegetation that touch the foundation. Rake away other potential pest harborage sites, such as deep mulch before application.
Firebrats		
Fleas		
Millipedes		
Mites		
Ticks		

PEST	RATE	USE INSTRUCTIONS
<p>Ants (Southern, Field, Allegheny Mound, Cornfield, Honey, Pavement, Nuisance, and Fire Ants) (excluding carpenter and pharaoh)</p> <p>Armyworms (Fall &amp; Yellow-striped)</p> <p>European Crane Flies</p> <p>Crickets</p> <p>Cutworms</p> <p>Earwigs</p> <p>Fleas</p> <p>Lawn moths (sod webworms)</p> <p>Grasshoppers</p> <p>Japanese beetle (adults)</p> <p>Spittlebugs</p> <p>Mites</p> <p>Green June Beetles</p> <p>Leafhoppers</p> <p>Millipedes</p> <p>Springtails</p> <p>Ticks</p>	<p>Apply at the rate of 2.2 lb to 3.3 lb per 1,000 sq. ft. (an area 20' x 50')</p> <p>or</p> <p>3.5 oz. to 5.3 oz. per 100 sq. ft. (10' x 10').</p>	<p>Apply with a fertilizer spreader (see spreader setting chart). Apply when grass is dry and water the lawn lightly immediately following application.</p> <p>Treat when insects first appear or when you notice lawn damage.</p>
<p>Billbugs</p> <p>Grubs (Japanese beetle grubs, Green June Beetle Grubs)</p> <p>Ataenius spretulus (dung beetles)</p> <p>Hyperodes Weevils</p> <p>Chinch bugs</p> <p>Mole Crickets</p>	<p>Apply 3.3 lb. per 1,000 sq. ft. of lawn.</p>	<p>Water grass completely immediately following application.</p> <p>Treat when insects first appear or when you notice lawn damage.</p> <p>For <b>hyperodes weevils</b>, apply in April-May.</p> <p>For <b>billbug control</b>, apply in May when activity is first observed or when chewed or brown grass indicates damage from these insects.</p> <p>For <b>grub control</b>, except on dung beetle, treat any time between late July and early October.</p> <p>For <b>dung beetle grub control</b>, apply in early spring or early summer (March-July) when damage is first noticed. Immediately following application, completely water grass with <math>\frac{1}{4}</math> to <math>\frac{1}{2}</math> inch of water. If a second generation occurs, treat again in late summer or early fall.</p> <p>For <b>chinch bug control</b>, apply during the summer when both nymphs and adults may be present.</p> <p>For <b>mole crickets</b> apply in the early spring for control of adults when actively invading grass areas. If the soil is dry, water before application to bring mole crickets closer to the surface and in contact with the granules. Apply late in the day and water in following application. For control of early nymphal stages, apply at the time of peak egg hatch. As the nymphs mature, higher rates may be necessary.</p>

<p>Ants (Southern, Field, Allegheny Mound, Cornfield, Honey, Pavement, Nuisance, and Fire Ants) (excluding carpenter and pharaoh)</p>	<p>Apply 1 teaspoon to ½ cup over, and 2 ft. around each ant mound</p>	<p>Use both broadcast and spot treatment for difficult to large ant infestations. If soil is dry, water lightly before application to bring the ants closer to the surface for better control. Immediately following application, water in completely. Apply a minimum of 1 gallon of water to each mound area. To prevent the disturbance of ants, water gently. Use of high pressure watering equipment may disturb the ants and cause migration, reducing product effectiveness. For best results, apply in cool weather, 65°-80°F, or in early morning or late evening hours. Treat new mounds as they appear.</p>
---	--	--

SUGGESTED SPREADER SETTINGS			
Spreader Model	LOW RATE 2.2 lbs/1,000 sq. ft.	MEDIUM RATE 3.4 lbs/1,000 sq. ft.	HIGH RATE 4.6 lbs/1,000 sq. ft.
The Andersons Yard Star 2150	13	14 1/2	15 1/2
The Andersons LCO 1000	K 1/4	L 1/2	M 1/2
Agri-Fab Broadcast (1-10)	4	5 1/2	6
Agri-Fab Drop (1-10)	4	5 1/2	6
Brinly 20 Series Push Broadcast (0-30)	13	14 1/2	15 1/2
Chapin Broadcast (1-30)	13	14 1/2	15 1/2
Craftsman Broadcast (0-30)	4	5 1/2	6
EarthWay Broadcast (0-30)	13	14 1/2	15 1/2
EarthWay Drop (0-30)	13	14 1/2	15 1/2
EarthWay 3400 Hand Spreader (1-3)	1	2	3
Precision Broadcast (1-10)	2 3/4	3 1/2	4 3/4
Precision Drop (1-10)	2 1/2	3	4 1/4
Scotts Broadcast (2-15) (includes Edgegard, Elite, Standard, Basic, LawnPro, & Speedy Green 1000-3000)	4	5 1/2	6
Scotts Drop (Accugreen 1000-2000 & Classic)	6	7 1/2	8
Scotts Wizz Hand Broadcast Spreader	4	5	6
Scotts Hand-Held Broadcast (Handy Green, Easy)	2	3	4
Spyker Broadcast	4.8	5.2	5.5

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in original container. Keep area inaccessible to children and pets.

**PESTICIDE DISPOSAL:** Pesticide wastes may be hazardous. Improper disposal of excess pesticide is a violation of Federal Law. Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

**CONTAINER HANDLING:** Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment, then offer for recycling if available, or dispose of empty bag in a sanitary landfill or by incineration.

The directions for use of this product must be followed carefully to the EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL, TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL, WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

Distributed by:  
The Andersons, Inc.  
P.O. Box 119  
Maumee, OH 43537  
[www.andersonsplantnutrient.com](http://www.andersonsplantnutrient.com)

GrubOut is a registered trademark of The Andersons.



The Andersons Professional with Grubout Plus (40lb)  
New

**10007802**  
F22

**Net Wt. 40 lb (18.14 kg)**