

More For Your Home™




Premium

Crabgrass Preventer plus Lawn Food

A herbicide plus fertilizer combination product that provides control of crabgrass and other weeds in established lawns and ornamental turfgrasses.

26•0•6

For Thick, Green, Beautiful Lawns



COVERS 6,000 SQUARE FEET



Net Wt. **17** Lbs. (7.71 kg)

- **Controls Crabgrass All Season Long**
- **Extended Greening From Controlled Release Nitrogen**

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See back panel for additional precautionary statements

EPA Reg. No. 9198-120
EPA Est. No. 9198-OH-1M, 9198-OH-2B, 9198-AL-001A
(Underlined letter used is first letter of run code on end of bag.)

ACTIVE INGREDIENT:

*Dithiopyr, 3,5-pyridinedicarbothioic acid, 2-(difluoromethyl)-4-(2-methylpropyl)-6-(trifluoromethyl)-S,S- dimethyl ester	0.103%
INERT INGREDIENTS	99.897%
TOTAL	100.00%

"More For Your Home" & The Andersons logo are registered trademarks of The Andersons, Inc.

More For Your Home™

Crabgrass Preventer
plus Lawn Food

The Andersons
Premium





Crabgrass Preventer

plus Lawn Food

GUARANTEED ANALYSIS

Total Nitrogen (N)	26.00%
26.00% Urea Nitrogen*	
Soluble Potash (K ₂ O)	6.00%
Derived from urea, potassium chloride, polymer coated sulfur coated urea.	
*3.90% Slowly Available Urea Nitrogen from Polymer Coated Sulfur Coated Urea	

F002091

Information regarding the contents and levels of metals in this product is available on the internet at <https://aapfco.org/metals>

Nitrogen Notice For those states with nitrogen application restrictions, including but not limited to MD, NH, NJ and VA, do not apply more than 2.8 lbs. of this product per 1,000 sq.ft.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

Keep Out of Reach of Children

CAUTION

Causes eye irritation. Harmful if inhaled. Avoid contact with eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Prolonged or frequently repeated skin contact while handling this material may cause allergic reaction in some individuals.

Directions For Use

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

New York Use Restrictions: Do not apply more than 5.6 pounds of this product per 1,000 square feet of turfgrass area per year. Do not make more than one application per year. Use only according to label instructions.

GENERAL INFORMATION

This product is a combination product that provides both crabgrass control and the required nutrients for optimum turfgrass growth all in a single product.

This product contains Dimension[®], a crabgrass preventer that can be applied up to 4 weeks later than other crabgrass preventers and provides up to 4 months control of crabgrass and other grass and broadleaf weeds prior to their establishment in lawns and ornamental turfgrasses. This product should be applied to lawns and ornamental turfgrasses that are well established (See "Use Precautions" section below for a list of the tolerant turfgrasses). This product will not harm nearby established ornamentals when used according to label directions.

USE PRECAUTIONS

This product should only be applied to lawns and ornamental turfgrasses that are well established. Use of this product on lawns and ornamental turfgrasses that are not well established, or have been weakened by stress due to mechanical operations, pests, excessive temperatures, or lack of moisture may increase the chances for turfgrass injury.

This product can be applied to the following turfgrasses when applied as directed:

Cool-Season Turfgrasses

Creeping Bentgrass
Fine Fescues
Kentucky Bluegrass
Perennial Ryegrass
Tall Fescues

Warm-Season Turfgrasses

Bahiagrass
Bermudagrass
Buffalograss
Carpetgrass
Centipedegrass
St. Augustinegrass
Zoysiagrass

This product should be applied in the spring or fall prior to the weed seed germination in established lawns and ornamental turfgrasses.

This product can be applied up to 4 weeks later than other crabgrass preventers and still provide season-long weed control of crabgrass. Dimension herbicide provides preemergence control of crabgrass and other weeds when applied prior to their germination, and it also provides postemergence control of crabgrass. The postemergence control provided by this product is limited to crabgrass only and will provide control of crabgrass when applied up to 4 weeks after crabgrass germination. Application of this product later than 4 weeks after crabgrass germination will result in reduced control of crabgrass.

For control of other grass and broadleaf weeds, this product must be applied as a preemergent treatment. Only crabgrass can be controlled with delayed applications after germination.

Tips for Improved Weed Control: For best results, apply this product within a few days after mowing and delay further mowings after the application of this product until the lawn or ornamental turfgrass has been watered or received significant rainfall to dissolve the fertilizer granules. Improved weed control may be observed when treated lawn or ornamental turfgrass is watered or receives rainfall within a few days after application of this product.

Reseeding, Overseeding or Sprigging of turfgrasses into areas that have been previously treated with this product should be delayed until 16 weeks after application. Reseeding, overseeding, or sprigging up to 16 weeks after application may prevent the establishment of newly reseeded, overseeded or sprigged turfgrasses.

- DO NOT apply more than 11 pounds of this product per 1,000 square feet of turfgrass area during the year (1,000 square feet area = 100 feet x 10 feet).
- DO NOT apply to flowers, vegetables, or fruit trees.
- DO NOT use clippings from treated turf for mulching around flowers, vegetables or fruit trees.
- Keep people and pets off treated areas until dust has settled.

APPLICATION DIRECTIONS

This product will provide control of crabgrass and other grasses and broadleaf weeds listed below when applied prior to weed seed germination in established lawns and ornamental turfgrasses.

Weeds Controlled and/or Suppressed by This Product

Grass Weeds:

Annual Bluegrass
Annual Ryegrass
Barnyardgrass
Crabgrass
Crowfootgrass
Goosegrass

Green Foxtail

Kikuyugrass (seedling)
Perennial Ryegrass
Smutgrass (seedling)
Yellow Foxtail

Broadleaf Weeds:

Bittercress
Buttercup Oxalis
Carolina Geranium
Chickweed
Common Lespedeza
Common Purslane
Corn Speedwell
Creeping Oxalis
Henbit
Parsley-plant
Pineappleweed
Spotted Spurge
Yellow Woodsorrel

Application Timings:

This product may be applied either in the spring or in the fall for the control of weeds.

When this product is applied in the fall, a sequential or second application should be made the following spring to provide season-long control of weeds.

Length of Control:


This product will provide up to 4 months of weed control if applied according to label directions.

The actual length of control will vary depending on use rate, weather conditions, condition of the turfgrass, intensity of weed pressure, and the location or "Region" within the United States that this product is being applied.

Please check the map to the right to determine which "Region" of the U.S. that applies to your particular use area.

Use "Regions" within the United States:

☐ North ☐ South



Recommended Use Rates

Recommended Use Rates for This Product (Pounds of this product to apply per 1,000 square feet of turfgrass)		
Region of U.S.	Length of Weed Control	
	2-3 Months Control	3-4 Months Control
North	2.8	4.0
South	4.3	5.6

FIRST AID If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contacts, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice. If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for treatment advice. If swallowed: Call a poison control center or doctor for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person. If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-757-8951 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. Use with care when applying to turf areas adjacent to any body of water. Drift and runoff from treated turf may adversely affect aquatic organisms in adjacent aquatic sites. Do not apply when weather conditions favor drift from treated areas. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

ANNUAL FERTILIZATION GUIDELINES FOR ESTABLISHED TURFGRASS LAWNS IN THREE REGIONS OF FLORIDA NITROGEN RECOMMENDATIONS (LBS N PER 1000 SQ. FT.)*					
Species	Bahiagrass	Bermuda	Centipede	St. Augustine	Zoysia
Timing of Application	North*/ Maximum Pounds N per Application (Only Apply to Actively Growing Turf)				
Spring or Summer	2	2	2	2	2
Fall	1	1	1	1	1
Maximum Annual Pounds	2-3	3-5	1-2	2-4	2-3
Timing of Application	Central*/ Maximum Pounds N per Application (Only Apply to Actively Growing Turf)				
Spring or Summer	2	2	2	2	2
Fall or Winter	1	1	1	1	1
Maximum Annual Pounds	2-4	4-6	2-3	2-5	2-4
Timing of Application	South*/ Maximum Pounds N per Application (Only Apply to Actively Growing Turf)				
Spring or Summer	2	2	2	2	2
Fall or Winter	1	1	1	1	1
Maximum Annual Pounds	2-4	5-7	2-3	4-6	2.5-4.5

*North Florida is defined as north of a line extending from Cedar Key to Daytona Beach. Central Florida is defined as south of the line extending from Cedar Key to Daytona Beach to a line extending from Tampa to Vero Beach. South Florida includes the remaining southern portion of the state.

STORAGE AND DISPOSAL


Do not contaminate water, food or feed by storage and disposal.
STORAGE: Keep container closed and away from food, feedstuffs and domestic water supplies.
DISPOSAL:
If empty: Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available.
If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

SUGGESTED SPREADER SETTINGS


IMPORTANT: It is important to follow the directions on this quality lawn care product as over-application may cause injury to the lawn. Measure the size of the area to be treated and do not over or under apply. Spreader settings are approximate. Differences may occur due to condition of spreader, speed at which applied and roughness of area treated. Calibrate spreader prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing uniformly. Close spreader when filling, stopping or turning. DO NOT HAND SPREAD.

BROADCAST SPREADER SETTINGS

For best results, apply in a circular path starting on the outside and working toward the center (see diagram below). A little overlap of product will ensure complete coverage.

	Spreader	2.8# Rate	4.0# Rate	4.3# Rate	5.6# Rate
		The Andersons Yard Star 2150 (0-30)13 1/2	15	15 1/2	17
	Agri-Fab Broadcast (1-10).....	4	5	5 1/4	6 1/2
	Brinly 20 Series Push Broadcast (0-30).....	13 1/2	15	15 1/2	17
	Chapin Broadcast (0-30).....	13 1/2	15	15 1/2	17
	Craftsman Broadcast (1-10).....	4	5	5 1/4	6 1/2
	EarthWay Broadcast (0-30).....	13 1/2	15	15 1/2	17
	Precision Broadcast (1-10).....	2 1/2	3 1/2	3 3/4	5
	Scotts Broadcast (Includes EdgeGuard, Elite, Standard, Basic, LawnPro, & Speed Green 1000-3000).....	4 1/2	5 1/2	5 3/4	7
	Spyker Broadcast.....	4.8	5.3	5.4	5.9

DROP SPREADER SETTINGS

	Spreader	2.8# Rate	4.0# Rate	4.3# Rate	5.6# Rate
		Agri-Fab Drop (1-10).....	5	5 1/4	6 1/2
	EarthWay Drop (0-30).....	13 1/2	15	15 1/2	17
	Precision Drop (1-10).....	2 1/2	3 1/2	3 3/4	5
	Scotts Drop (Accugreen 1000-3000 & Classic).....	6 1/4	7 1/2	7 3/4	9

IMPORTANT: READ BEFORE USE:

Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user of Buyer accepts the following Warranty Disclaimer and Limitation of Liability.

WARRANTY DISCLAIMER and LIMITATION of LIABILITY:

Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions. Manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. To the extent consistent with applicable law, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyer's choice. To the extent consistent with applicable law, Manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted.

LEGAL RIGHTS:

THE LIMITED WARRANTY GIVES YOUR SPECIFIC LEGAL RIGHTS; YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATE/JURISDICTION TO STATE/JURISDICTION.

Manufactured by:
The Andersons, Inc.
P.O. Box 119
Maumee, OH 43537

031865 32056 9
FPO

AND263DM1
AND263DM1B
©2021

082421