

More For Your Home™



Premium

Weed & Feed

24•0•16

For Thick,
Green Lawns

COVERS 5,000 SQUARE FEET



Net Wt. **18** Lbs. (8.16 kg)

- **Controls Dandelions
& Over 250 Other
Broadleaf Weeds**

- **A Premium Blend of
Fertilizer Nutrients
that Provide Quick &
Extended Greening**

- **Contains the 3-way
Power of Dissolve®
for Broadleaf,
Creeping & Woody
Weeds**



**KEEP OUT OF REACH OF CHILDREN
CAUTION**

SEE BACK PANEL FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800)424-9300.

For Medical Emergencies Only, Call (877) 325-1840.

EPA Reg. No. 228-281-9198

EPA Est. No. 9198-OH-1M, 9198-OH-2B, 9198-AL-001A

(Underlined letter used is first letter on run code on end of bag.)

ACTIVE INGREDIENTS:

Dimethylamine Salt of 2,4-Dichlorophenoxyacetic Acid*	0.560%
Dimethylamine Salt of (+)-R-2-(2-Methyl-4-Chlorophenoxy)propionic Acid**†	0.145%
Dimethylamine Salt of (+)-R-2-(2,4-Dichlorophenoxy)propionic Acid***†	0.143%
OTHER INGREDIENTS:	99.152%
TOTAL	100.000%

ISOMER SPECIFIC METHOD, EQUIVALENT TO:

*2,4-Dichlorophenoxyacetic Acid	0.46%
**(+)-R-2-(2-Methyl-4-Chlorophenoxy)propionic Acid	0.12%
***(+)-R-2-(2,4-Dichlorophenoxy)propionic Acid	0.12%

†CONTAINS THE SINGLE ISOMER FORMS OF MECOPROP-p AND DICHLORPROP-p.

"More For Your Home" & The Andersons logo are registered trademarks of The Andersons, Inc.;
Dissolve is a registered trademark of Nufarm Americas, Inc.



More For Your Home™



Premium

Weed & Feed



Weed & Feed

GUARANTEED ANALYSIS

Total Nitrogen (N) 24.0%
24% Urea Nitrogen*
Soluble Potash (K₂O) 16.0%
Iron (Fe) 1.0%
Derived from urea, potassium chloride, polymer coated sulfur coated urea and ferric oxide.
***7.0% Slowly Available Nitrogen from Polymer Coated Sulfur Coated Urea F00292**
Fertilizer Guaranteed by: The Andersons Lawn Fertilizer Division, Inc., P. O. Box 119, Maumee, OH 43537

A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed, this product will supplement your regular fertilizer program by providing 0.86 pounds of nitrogen per 1,000 square feet and herbicide to control listed weeds. Please consult your local extension office for questions regarding a regular fertilizer program.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Causes moderate eye irritation. Avoid contact with eyes or clothing. Harmful if inhaled or absorbed through skin. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

	FIRST AID
If in eyes:	<ul style="list-style-type: none">Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.Call a poison control center or doctor for treatment advice.
If on skin or clothing:	<ul style="list-style-type: none">Take off contaminated clothing.Rinse skin immediately with plenty of water for 15 to 20 minutes.Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none">Call a poison control center or doctor immediately fir treatment advice.Have person sip a glass of water if able to swallow.Do not induce vomiting unless told to do so by a poison control center or doctor.Do not give anything by mouth to an unconscious person.
HOTLINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.	

ENVIRONMENTAL HAZARDS

This product is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

DIRECTIONS FOR USE

It is violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled. Do not apply more than 2 broadcast applications per treatment site per year with minimum of 30 days between applications.

This product is a stable mixture of three herbicides. It is for use on all Residential Turf.

This product may be used on Bahia, common Bermuda, Bluegrass, Centipede, Fescue, Rye, Rye, St. Augustine and Zoysia grass. DO NOT USE on Dichondra and Carpet grass. DO NOT USE this product where desirable clovers are present or allow the product to “wind drift” onto flowers, vegetables, ornamentals, shrubs, trees or other desirable plants.

WHEN TO APPLY

Best results are obtained when weeds are growing actively in Spring or early Fall. Avoid contact with desirable shrubs, plants, vegetables, or flowers; especially when applying with a rotary (spinner) type spreader. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded or sodded lawns until after 3 mowings. For optimum results (1) avoid mowing 1 to 2 days before and after application; (2) apply when grass is moist (after a rain, sprinkling or dew); (3) avoid unnecessary disturbances, including watering of treated areas for 48 hours after application. Watering will wash off weed killing material; and (4) avoid applying if rainfall is expected within 48 hours following treatment. Possible retreatment in approximately 30 days may be necessary if lawn if heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed recommended rates because damage to turf may occur.

NOTE: When applying to Southern grasses - under conditions of inadequate moisture, high temperature and high humidity, over-application may cause minor transitory discoloration on Centipede and St. Augustine grasses. It is recommended that during hot, dry conditions, application rates should be reduced 50% to avoid possible dehydration or browning of grass.

HOWTO APPLY

This 18 pound bag should be used on 5,000 square feet. Apply evenly with lawn spreader as a one pass application. A two pass application, criss-crossing the area at half the rate each time is equal to one application.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Store in original container and keep any bags containing unused product in locked storage inaccessible to children and domestic animals. Do not store near fees or foodstuffs. To prevent cross-contamination, do not store near open containers of fertilizer, seed, or other pesticides.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. **If empty** - Place in trash or offer for recycling if available. **If party filled** - Call your local solid waste agency for disposal instructions. Never place unused product down any indoor (inducing toilet) or outdoor (including sewer) drain.

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOUR ARE FURNISHED “AS IS” BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR Y USAGE OF TRADE, STATUTORY O OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OR MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THERE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR RISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER,S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded. (RV070609)

This product is designed to promoted lush green growth in lawns and control many broadleaf weeds such as:

Alder	Cinquefoil	Galinsoga, Smallflower	Oxalis	Spurweed
Artichoke	Cinquefoil, Rough	Geranium, Carolina	Pansy, Field	Stinkweed
Aster	Clematis, Western	Goathead	Plantain, Blackseed	Stitchwort
Austrian Fieldcress	Clover	Goatsbeard	Parsnip	Sumac
Bedstraw	Clover, Annual	Goldenrod	Pearlwort	Sunflower
Bedstraw, Pineywoods	Yellow Sweet	Ground Ivy	Pennycress	Sweet Clover
Beggartick	Clover, Crimson	Groundsel	Pennywort	Tansy Ragwort
Betony, Florida	Clover, Hop	Gumweed	Peppergrass	Tanweed
Biden	Clover, Red	Hawkweed	Pepperweed	Tarweed
Bindweed	Clover, Strawberry	Healall	Pigweed	Thistle
Bindweed, Field	Clover, Sweet	Heartleaf Drymary	Plantain	Toadflax
Bindweed, Hedge	Clover, White	Hemlock, Poison	Poison Hemlock	Tumbleweed
Bitter wintercress	Cockle	Hemp	Poison Ivy	Velvet Leaf
Bittercress, Hairy	Cocklebur	Henbit	Poison Oak	Veronica
Bitterweed	Cocklebur, Spiny	Hoary Cress	Pokeweed	Vervain
Black-eyed Susan	Coffeebean	Honeysuckle	Poorjoe	Vetch
Black Medic	Coffeeweed	Horsetail	Povertyweed	Virginia
Blessed Thistle	Common Sowthistle	Indiana Mallow	Prickly Lettuce	Buttonweed
Bloodflower, Milkweed	Coreopsis, Plains	Ironweed	Prickly Sida	Virginia Creeper
Blue Lettuce	Creeping Jenny	Jewelweed	Primrose	Wild Aster
Box Elder	Croton	Jimsonweed	Puncture Vine	Wild Carrot
Brassbuttons	Croton, Wooly	Kochia	Purslane	Wild Garlic
Bristly Oxtongue	Cudweed	Knapweed, Spotted	Ragweed	Wild Geranium
Broomweed	Curly Indigo	Knawel	Red Clover	Wild Lettuce
Buckhorn	Daisy, Oweye	Knotweed	Red Sorrel	Wild Onion
Buckwheat, Wild	Dandelion	Knotweed, Prostrate	Rush	Wild Radish
Bull Thistle	Dock	Lambsquarter	Russian Thistle	Wild Rape
Burdock	Dogbane	Lespedeza	St. Johnswort	Wild Strawberry
Bur Ragweed	Dogfennel	Locoweed	Salsify, Western	Wild Sweet
Burweed	Elderberry	Lupines	Scarlet Pimpernel	Potato
Buttercup	English Daisy	Mallow	Sheep Sorrel	Willow
Buttercup, Bulbous	Fiddleneck	Mallow, Venice	Shepherdspurse	Witchweed
Canada Thistle	Filaree, Redstem	Marigold, Wild	Smartweed	Woodsorrel
Carpetweed	Flax, False	Marshelder	Sneezeweed	Wormseed
Catchweed Bedstraw	Flea Bane (Daisy)	Matchweed	SouthernWild	Yarrow
Catnip	Fleabane, Hairy	Mexicanweed	Rose	Yellow Rocket
Catsear	Fleabane, Rough	Morningglory	Sowthistle	Yellowflower
Catsear, Spotted	Flixweed	Mullein, Common	Spanishneedle	Pepperweed
Chamomile, Corn	Florida Pusley	Musk Thistle	Spatterdock	and other
Chickweed	Four-O-Clock, Wild	Mustard	Speedwell	broadleaf weeds
Chickweed, Common	Frenchweed	Nettle	Spiny Amaranth	
Chicory	Galinsoga	Orange Hawkweed	Spurge	

The following settings are **approximate** for applying contents of this package at a suggested rate of 3.6 pounds per 1,000 square feet. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed. Check your spreader on 250 square feet and adjust to apply at the rate of 0.9 pound per 250 square feet.

BROADCAST SPREADER SETTINGS

FOR ONE PASS APPLICATION: For best results, apply in a circular path starting on the outside and working toward the center (see broadcast spreader path below). A little overlap of product will ensure complete coverage.


	3.6# Rate
The Andersons 4300	4
The Andersons 4500	5 1/2
Precision 4300	4
Precision 4500	5 1/2
Republic EZ 540/75450	2 - 2 1/2
Scotts Basic, Scotts Standard . .	3 1/2 - 4
Scotts Lawn Pro, Scotts Deluxe .	3 1/2 - 4
Scotts SpeedyGreen SG3000 . .	3 1/2 - 4
Scotts EasyGreen EG3	24 1/2
Scotts Edgeguard	4



DROP SPREADER SETTINGS

FOR ONE PASS APPLICATION: First apply 2 border strips along the edge of each end of the lawn (see drop spreader path diagram below). Then apply product along the length of the lawn, overlapping slightly and shutting off the spreader before turning as you reach the edge of the border strips.

	3.6# Rate
Precision DS4500	2 1/2
Republic EZ 530/75350	5 - 5 1/2
Scotts PF3	4 - 4 1/2
Scotts Accugreen	6



Sweep (do not wash) all fertilizer materials from surfaces immediately after application, because certain fertilizer ingredients can cause permanent stains. Wash out spreader after each use and allow it to dry thoroughly. Treat any exposed metal spreader areas with oil to reduce corrosion.

Distributed by:
The Andersons, Inc.
P. O. Box 119
Maumee, OH 43537

©2020

AND247DSFE18
AND247DSFE18B

